

ON NEIGHBORHOOD AND PARTIAL SUMS PROBLEM FOR GENERALIZED SAKAGUCHI TYPE FUNCTIONS

MURAT AĞLAR AND HALIT ORHAN

ABSTRACT. In the present investigation, we introduce a new class $k - \widetilde{\mathcal{US}}_s^n(\lambda, \mu, \gamma, t)$ of analytic functions in the open unit disc $\mathcal{U}$ with negative coefficients. The object of the present paper is to determine coefficient estimates, neighborhoods and partial sums for functions $f(z)$ belonging to this class.

1. Introduction

Let $\mathcal{A}$ denote the family of functions f of the form

$$(1.1) \quad f(z) = z + \sum_{n=2}^{\infty} a_n z^n$$

that are analytic in the open unit disk $\mathcal{U} = \{z : |z| < 1\}$. Denote by $\mathcal{S}$ the subclass of $\mathcal{A}$ of functions that are univalent in $\mathcal{U}$.

For $f \in \mathcal{A}$ given by (1.1) and $g(z)$ given by

$$(1.2) \quad g(z) = z + \sum_{n=2}^{\infty} b_n z^n$$

their convolution (or Hadamard product), denoted by $(f * g)$, is defined as

$$(1.3) \quad (f * g)(z) := z + \sum_{n=2}^{\infty} a_n b_n z^n =: (g * f)(z) \quad (z \in \mathcal{U}).$$

Note that $f * g \in \mathcal{A}$.

A function $f \in \mathcal{A}$ is said to be in $k - \mathcal{US}(\gamma)$, the class of k -uniformly starlike functions of order γ , $0 \leq \gamma < 1$, if satisfies the condition

$$(1.4) \quad \operatorname{Re} \left\{ \frac{z f'(z)}{f(z)} \right\} > k \left| \frac{z f'(z)}{f(z)} - 1 \right| + \gamma \quad (k \geq 0),$$

2000 *Mathematics Subject Classification.* 30C45.

Key words and phrases. Analytic function, uniformly starlike function, coefficient estimate, neighborhood problem, partial sums.

and a function $f \in \mathcal{A}$ is said to be in $k-\mathcal{UC}(\gamma)$, the class of k -uniformly convex functions of order γ , $0 \leq \gamma < 1$, if satisfies the condition

$$(1.5) \quad \operatorname{Re} \left\{ 1 + \frac{zf''(z)}{f'(z)} \right\} > k \left| \frac{zf''(z)}{f'(z)} \right| + \gamma \quad (k \geq 0).$$

Uniformly starlike and uniformly convex functions were first introduced by Goodman [8] and then studied by various authors. It is known that $f \in k-\mathcal{UC}(\gamma)$ or $f \in k-\mathcal{US}(\gamma)$ if and only if $1 + \frac{zf''(z)}{f'(z)}$ or $\frac{zf'(z)}{f(z)}$, respectively, takes all the values in the conic domain $\mathcal{R}_{k,\gamma}$ which is included in the right half plane given by

$$(1.6) \quad \mathcal{R}_{k,\gamma} := \left\{ w = u + iv \in \mathbb{C} : u > k\sqrt{(u-1)^2 + v^2} + \gamma, \beta \geq 0 \text{ and } \gamma \in [0, 1) \right\}.$$

Denote by $\mathcal{P}(P_{k,\gamma})$, ($\beta \geq 0$, $0 \leq \gamma < 1$) the family of functions p , such that $p \in \mathcal{P}$, where $\mathcal{P}$ denotes well-known class of Caratheodory functions. The function $P_{k,\gamma}$ maps the unit disk conformally onto the domain $\mathcal{R}_{k,\gamma}$ such that $1 \in \mathcal{R}_{k,\gamma}$ and $\partial\mathcal{R}_{k,\gamma}$ is a curve defined by the equality

$$(1.7) \quad \partial\mathcal{R}_{k,\gamma} := \left\{ w = u + iv \in \mathbb{C} : u^2 = \left(k\sqrt{(u-1)^2 + v^2} + \gamma \right)^2, \beta \geq 0 \text{ and } \gamma \in [0, 1) \right\}.$$

From elementary computations we see that (1.7) represents conic sections symmetric about the real axis. Thus $\mathcal{R}_{k,\gamma}$ is an elliptic domain for $k > 1$, a parabolic domain for $k = 1$, a hyperbolic domain for $0 < k < 1$ and the right half plane $u > \gamma$, for $\beta = 0$.

In [13], Sakaguchi defined the class $\mathcal{S}_s$ of starlike functions with respect to symmetric points as follows:

Let $f \in \mathcal{A}$. Then f is said to be starlike with respect to symmetric points in $\mathcal{U}$ if and only if

$$\operatorname{Re} \left\{ \frac{2zf'(z)}{f(z) - f(-z)} \right\} > 0 \quad (z \in \mathcal{U}).$$

Recently, Owa et. al. [10] defined the class $\mathcal{S}_s(\alpha, t)$ as follows:

$$\operatorname{Re} \left\{ \frac{(1-t)zf'(z)}{f(z) - f(tz)} \right\} > \alpha \quad (z \in \mathcal{U}),$$

where $0 \leq \alpha < 1$, $|t| \leq 1$, $t \neq 1$. Note that $\mathcal{S}_s(0, -1) = \mathcal{S}_s$ and $\mathcal{S}_s(\alpha, -1) = \mathcal{S}_s(\alpha)$ is called Sakaguchi function of order α .

The *linear multiplier differential operator* $D_{\lambda,\mu}^\eta f$ was defined by the authors in (see [11]) as follows

$$\begin{aligned} D_{\lambda,\mu}^0 f(z) &= f(z) \\ D_{\lambda,\mu}^1 f(z) &= D_{\lambda,\mu} f(z) = \lambda \mu z^2 (f(z))'' + (\lambda - \mu) z (f(z))' + (1 - \lambda + \mu) f(z) \\ D_{\lambda,\mu}^2 f(z) &= D_{\lambda,\mu} (D_{\lambda,\mu}^1 f(z)) \\ &\vdots \\ D_{\lambda,\mu}^\eta f(z) &= D_{\lambda,\mu} (D_{\lambda,\mu}^{\eta-1} f(z)) \end{aligned}$$

where $0 \leq \mu \leq \lambda \leq 1$ and $\eta \in \mathbb{N}_0 = \mathbb{N} \cup \{0\}$. Later, the operator $D_{\lambda,\mu}^\eta f$ was extended for $\lambda \geq \mu \geq 0$ by the authors in (see [5]).

If f is given by (1.1) then from the definition of the operator $D_{\lambda,\mu}^\eta f(z)$ it is easy to see that

$$(1.8) \quad D_{\lambda,\mu}^\eta f(z) = z + \sum_{n=2}^{\infty} \Phi^\eta(\lambda, \mu, n) a_n z^n$$

where

$$(1.9) \quad \Phi^\eta(\lambda, \mu, n) = [1 + (\lambda \mu n + \lambda - \mu)(n-1)]^\eta.$$

It should be remarked that the operator $D_{\lambda,\mu}^\eta$ is a generalization of many other linear operators considered earlier. In particular, for $f \in \mathcal{A}$ we have the following:

- $D_{1,0}^\eta f(z) \equiv D^\eta f(z)$ the operator investigated by Sălăgean (see [14]).
- $D_{\lambda,0}^\eta f(z) \equiv D_\lambda^\eta f(z)$ the operator studied by Al-Oboudi (see [1]).

Now, by making use of the differential operator $D_{\lambda,\mu}^\eta$, we define a new subclass of functions belonging to the class $\mathcal{A}$.

Definition 1.1. A function $f(z) \in \mathcal{A}$ is said to be in the class $k - \mathcal{US}_s^\eta(\lambda, \mu, \gamma, t)$ if for all $z \in \mathcal{U}$,

$$\operatorname{Re} \left\{ \frac{(1-t)z \left(D_{\lambda,\mu}^\eta f(z) \right)'}{D_{\lambda,\mu}^\eta f(z) - D_{\lambda,\mu}^\eta f(tz)} \right\} \geq k \left| \frac{(1-t)z \left(D_{\lambda,\mu}^\eta f(z) \right)'}{D_{\lambda,\mu}^\eta f(z) - D_{\lambda,\mu}^\eta f(tz)} - 1 \right| + \gamma$$

for $\lambda \geq \mu \geq 0$, $\eta, k \geq 0$, $|t| \leq 1$, $t \neq 1$, $0 \leq \gamma < 1$.

Furthermore, we say that a function $f(z) \in k - \mathcal{US}_s^\eta(\lambda, \mu, \gamma, t)$ is in the subclass $k - \widetilde{\mathcal{US}}_s^\eta(\lambda, \mu, \gamma, t)$ if $f(z)$ is of the following form:

$$(1.10) \quad f(z) = z - \sum_{n=2}^{\infty} a_n z^n \quad (a_n \geq 0, n \in \mathbb{N}).$$

The aim of the present paper is to study the coefficient bounds, partial sums and certain neighborhood results of the class $k - \widetilde{\mathcal{US}}_s^\eta(\lambda, \mu, \gamma, t)$.

Remark 1.1. *Through our present investigation, we tacitly assume that the parametric constraints listed (1.9).*

2. Coefficient bounds of the function class $k - \widetilde{\mathcal{US}}_s^\eta(\lambda, \mu, \gamma, t)$

Firstly, we shall need the following lemmas.

Lemma 2.1. *Let $w = u + iv$. Then*

$$\operatorname{Re} w \geq \alpha \quad \text{if and only if} \quad |w - (1 + \alpha)| \leq |w + (1 - \alpha)|.$$

Lemma 2.2. *Let $w = u + iv$ and α, γ are real numbers. Then*

$$\operatorname{Re} w > \alpha |w - 1| + \gamma \quad \text{if and only if} \quad \operatorname{Re} \{w(1 + \alpha e^{i\theta}) - \alpha e^{i\theta}\} > \gamma.$$

Theorem 2.1. *The function $f(z)$ defined by (1.10) is in the class $k - \widetilde{\mathcal{US}}_s^\eta(\lambda, \mu, \gamma, t)$ if and only if*

$$(2.1) \quad \sum_{n=2}^{\infty} \Phi^\eta(\lambda, \mu, n) |n(k+1) - u_n(k+\gamma)| a_n \leq 1 - \gamma,$$

where $\lambda \geq \mu \geq 0$, $\eta, k \geq 0$, $|t| \leq 1$, $t \neq 1$, $0 \leq \gamma < 1$, $u_n = 1 + t + \dots + t^{n-1}$.

The result is sharp for the function $f(z)$ given by

$$f(z) = z - \frac{1 - \gamma}{\Phi^\eta(\lambda, \mu, n) |n(k+1) - u_n(k+\gamma)|} z^n.$$

Proof. By Definition 1.1, we get

$$\operatorname{Re} \left\{ \frac{(1-t)z \left(D_{\lambda, \mu}^\eta f(z) \right)'}{D_{\lambda, \mu}^\eta f(z) - D_{\lambda, \mu}^\eta f(tz)} \right\} \geq k \left| \frac{(1-t)z \left(D_{\lambda, \mu}^\eta f(z) \right)'}{D_{\lambda, \mu}^\eta f(z) - D_{\lambda, \mu}^\eta f(tz)} - 1 \right| + \gamma.$$

Then by Lemma 2.2, we have

$$\operatorname{Re} \left\{ \frac{(1-t)z \left(D_{\lambda, \mu}^\eta f(z) \right)'}{D_{\lambda, \mu}^\eta f(z) - D_{\lambda, \mu}^\eta f(tz)} (1 + ke^{i\theta}) - ke^{i\theta} \right\} \geq \gamma, \quad -\pi < \theta \leq \pi$$

or equivalently

$$(2.2) \quad \operatorname{Re} \left\{ \frac{(1-t)z \left(D_{\lambda, \mu}^\eta f(z) \right)' (1 + ke^{i\theta})}{D_{\lambda, \mu}^\eta f(z) - D_{\lambda, \mu}^\eta f(tz)} - \frac{ke^{i\theta} [D_{\lambda, \mu}^\eta f(z) - D_{\lambda, \mu}^\eta f(tz)]}{D_{\lambda, \mu}^\eta f(z) - D_{\lambda, \mu}^\eta f(tz)} \right\} \geq \gamma.$$

Let

$$F(z) = (1-t)z \left(D_{\lambda, \mu}^\eta f(z) \right)' (1 + ke^{i\theta}) - ke^{i\theta} [D_{\lambda, \mu}^\eta f(z) - D_{\lambda, \mu}^\eta f(tz)]$$

and

$$E(z) = D_{\lambda, \mu}^{\eta} f(z) - D_{\lambda, \mu}^{\eta} f(tz).$$

By Lemma 2.1, (2.2) is equivalent to

$$|F(z) + (1 - \gamma)E(z)| \geq |F(z) - (1 + \gamma)E(z)| \quad \text{for } 0 \leq \gamma < 1.$$

But

$$\begin{aligned} |F(z) + (1 - \gamma)E(z)| &= \left| (1 - t) \left\{ (2 - \gamma)z - \sum_{n=2}^{\infty} \Phi^{\eta}(\lambda, \mu, n) (n + u_n(1 - \gamma)) a_n z^n \right. \right. \\ &\quad \left. \left. - k e^{i\theta} \sum_{n=2}^{\infty} \Phi^{\eta}(\lambda, \mu, n) (n - u_n) a_n z^n \right\} \right| \\ &\geq |1 - t| \left\{ (2 - \gamma)|z| - \sum_{n=2}^{\infty} \Phi^{\eta}(\lambda, \mu, n) |n + u_n(1 - \gamma)| a_n |z|^n \right. \\ &\quad \left. - k \sum_{n=2}^{\infty} \Phi^{\eta}(\lambda, \mu, n) |n - u_n| a_n |z|^n \right\}. \end{aligned}$$

Also

$$\begin{aligned} |F(z) - (1 + \gamma)E(z)| &= \left| (1 - t) \left\{ -\gamma z - \sum_{n=2}^{\infty} \Phi^{\eta}(\lambda, \mu, n) (n - (1 + \gamma)u_n) a_n z^n \right. \right. \\ &\quad \left. \left. - k e^{i\theta} \sum_{n=2}^{\infty} \Phi^{\eta}(\lambda, \mu, n) (n - u_n) a_n z^n \right\} \right| \\ &\leq |1 - t| \left\{ \gamma|z| + \sum_{n=2}^{\infty} \Phi^{\eta}(\lambda, \mu, n) |n - u_n(1 + \gamma)| a_n |z|^n \right. \\ &\quad \left. + k \sum_{n=2}^{\infty} \Phi^{\eta}(\lambda, \mu, n) |n - u_n| a_n |z|^n \right\} \end{aligned}$$

and so

$$\begin{aligned} &|F(z) + (1 - \gamma)E(z)| - |F(z) - (1 + \gamma)E(z)| \\ &\geq |1 - t| \left\{ 2(1 - \gamma)|z| - \sum_{n=2}^{\infty} \Phi^{\eta}(\lambda, \mu, n) [|n + u_n(1 - \gamma)| + |n - u_n(1 + \gamma)| + 2k|n - u_n|] a_n |z|^n \right\} \\ &\geq 2(1 - \gamma)|z| - \sum_{n=2}^{\infty} 2\Phi^{\eta}(\lambda, \mu, n) |n(k + 1) - u_n(k + \gamma)| a_n |z|^n \geq 0 \end{aligned}$$

or

$$\sum_{n=2}^{\infty} \Phi^{\eta}(\lambda, \mu, n) |n(k + 1) - (k + \gamma)u_n| a_n \leq 1 - \gamma.$$

Conversely, suppose that (2.1) holds. Then we must show

$$\operatorname{Re} \left\{ \frac{(1 - t)z \left(D_{\lambda, \mu}^{\eta} f(z) \right)' (1 + k e^{i\theta}) - k e^{i\theta} [D_{\lambda, \mu}^{\eta} f(z) - D_{\lambda, \mu}^{\eta} f(tz)]}{D_{\lambda, \mu}^{\eta} f(z) - D_{\lambda, \mu}^{\eta} f(tz)} \right\} \geq \gamma.$$

Upon choosing the values of z on the positive real axis where $0 \leq z = r < 1$, the above inequality reduces to

$$\operatorname{Re} \left\{ \frac{(1 - \gamma) - \sum_{n=2}^{\infty} \Phi^{\eta}(\lambda, \mu, n) (n(1 + ke^{i\theta}) - u_n(\gamma + ke^{i\theta})) a_n z^{n-1}}{1 - \sum_{n=2}^{\infty} \Phi^{\eta}(\lambda, \mu, n) u_n a_n z^{n-1}} \right\} \geq 0.$$

Since $\operatorname{Re}(-e^{i\theta}) \geq -|e^{i\theta}| = -1$, the above inequality reduces to

$$\operatorname{Re} \left\{ \frac{(1 - \gamma) - \sum_{n=2}^{\infty} \Phi^{\eta}(\lambda, \mu, n) (n(1 + k) - u_n(\gamma + k)) a_n r^{n-1}}{1 - \sum_{n=2}^{\infty} \Phi^{\eta}(\lambda, \mu, n) u_n a_n r^{n-1}} \right\} \geq 0.$$

Letting $r \rightarrow 1^-$, we have desired conclusion. $\square$

Corollary 2.2. *If $f(z) \in k - \widetilde{\mathcal{US}}_s^{\eta}(\lambda, \mu, \gamma, t)$, then*

$$a_n \leq \frac{1 - \gamma}{\Phi^{\eta}(\lambda, \mu, n) |n(k + 1) - u_n(k + \gamma)|}$$

where $\lambda \geq \mu \geq 0$, $\eta, k \geq 0$, $|t| \leq 1$, $t \neq 1$, $0 \leq \gamma < 1$, $u_n = 1 + t + \dots + t^{n-1}$.

3. Neighborhood of the function class $k - \widetilde{\mathcal{US}}_s^{\eta}(\lambda, \mu, \gamma, t)$

Following the earlier investigations (based upon the familiar concept of neighborhoods of analytic functions) by Goodman [7], Ruscheweyh [12], Altıntaş *et al.* ([2], [3]) and others including Srivastava *et al.* ([17], [18]), Orhan ([9]), Deniz *et al.* [6], Cataş [4].

Definition 3.1. *Let $\lambda \geq \mu \geq 0$, $\eta, k \geq 0$, $|t| \leq 1$, $t \neq 1$, $0 \leq \gamma < 1$, $\alpha \geq 0$, $u_n = 1 + t + \dots + t^{n-1}$.*

We define the α -neighborhood of a function $f \in \mathcal{A}$ and denote by $\mathcal{N}_{\alpha}(f)$ consisting of all functions

$g(z) = z - \sum_{n=2}^{\infty} b_n z^n \in \mathcal{S}$ ($b_n \geq 0$, $n \in \mathbb{N}$) satisfying

$$\sum_{n=2}^{\infty} \frac{\Phi^{\eta}(\lambda, \mu, n) |n(k + 1) - u_n(k + \gamma)|}{1 - \gamma} |a_n - b_n| \leq \alpha.$$

Theorem 3.1. *Let $f \in k - \widetilde{\mathcal{US}}_s^{\eta}(\lambda, \mu, \gamma, t)$ and for all real θ we have $\gamma(e^{i\theta} - 1) - 2e^{i\theta} \neq 0$. For any complex number ε with $|\varepsilon| < \alpha$ ($\alpha \geq 0$), if f satisfies the following condition:*

$$\frac{f(z) + \varepsilon z}{1 + \varepsilon} \in k - \widetilde{\mathcal{US}}_s^{\eta}(\lambda, \mu, \gamma, t),$$

then $\mathcal{N}_{\alpha}(f) \subset k - \widetilde{\mathcal{US}}_s^{\eta}(\lambda, \mu, \gamma, t)$.

Proof. It is obvious that $f \in k - \widetilde{\mathcal{US}}_s^\eta(\lambda, \mu, \gamma, t)$ if and only if

$$\left| \frac{(1-t)z \left(D_{\lambda, \mu}^\eta f(z) \right)' (1 + ke^{i\theta}) - (ke^{i\theta} + 1 + \gamma) \left(D_{\lambda, \mu}^\eta f(z) - D_{\lambda, \mu}^\eta f(tz) \right)}{(1-t)z \left(D_{\lambda, \mu}^\eta f(z) \right)' (1 + ke^{i\theta}) + (1 - ke^{i\theta} - \gamma) \left(D_{\lambda, \mu}^\eta f(z) - D_{\lambda, \mu}^\eta f(tz) \right)} \right| < 1 \quad (-\pi < \theta < \pi)$$

for any complex number s with $|s| = 1$, we have

$$\frac{(1-t)z \left(D_{\lambda, \mu}^\eta f(z) \right)' (1 + ke^{i\theta}) - (ke^{i\theta} + 1 + \gamma) \left(D_{\lambda, \mu}^\eta f(z) - D_{\lambda, \mu}^\eta f(tz) \right)}{(1-t)z \left(D_{\lambda, \mu}^\eta f(z) \right)' (1 + ke^{i\theta}) + (1 - ke^{i\theta} - \gamma) \left(D_{\lambda, \mu}^\eta f(z) - D_{\lambda, \mu}^\eta f(tz) \right)} \neq s.$$

In other words, we must have

$$(1-s)(1-t)z \left(D_{\lambda, \mu}^\eta f(z) \right)' (1 + ke^{i\theta}) - (ke^{i\theta} + 1 + \gamma + s(ke^{i\theta} - 1 + \gamma)) \left(D_{\lambda, \mu}^\eta f(z) - D_{\lambda, \mu}^\eta f(tz) \right) \neq 0$$

which is equivalent to

$$z - \sum_{n=2}^{\infty} \frac{\Phi^\eta(\lambda, \mu, n) \left((n - u_n)(1 + ke^{i\theta} - ske^{i\theta}) - s(n + u_n) - u_n\gamma(1 - s) \right)}{\gamma(s - 1) - 2s} z^n \neq 0.$$

However, $f \in k - \widetilde{\mathcal{US}}_s^\eta(\lambda, \mu, \gamma, t)$ if and only if $\frac{(f*h)(z)}{z} \neq 0$, $z \in \mathcal{U} - \{0\}$ where $h(z) = z - \sum_{n=2}^{\infty} c_n z^n$,

and

$$c_n = \frac{\Phi^\eta(\lambda, \mu, n) \left((n - u_n)(1 + ke^{i\theta} - ske^{i\theta}) - s(n + u_n) - u_n\gamma(1 - s) \right)}{\gamma(s - 1) - 2s}$$

we note that

$$|c_n| \leq \frac{\Phi^\eta(\lambda, \mu, n) |n(1 + k) - u_n(k + \gamma)|}{1 - \gamma}$$

since $\frac{f(z) + \varepsilon z}{1 + \varepsilon} \in k - \widetilde{\mathcal{US}}_s^\eta(\lambda, \mu, \gamma, t)$, therefore $z^{-1} \left(\frac{f(z) + \varepsilon z}{1 + \varepsilon} * h(z) \right) \neq 0$, which is equivalent to

$$(3.1) \quad \frac{(f * h)(z)}{(1 + \varepsilon)z} + \frac{\varepsilon}{1 + \varepsilon} \neq 0.$$

Now suppose that $\left| \frac{(f*h)(z)}{z} \right| < \alpha$. Then by (3.1), we must have

$$\left| \frac{(f * h)(z)}{(1 + \varepsilon)z} + \frac{\varepsilon}{1 + \varepsilon} \right| \geq \frac{|\varepsilon|}{|1 + \varepsilon|} - \frac{1}{|1 + \varepsilon|} \left| \frac{(f * h)(z)}{z} \right| > \frac{|\varepsilon| - \alpha}{|1 + \varepsilon|} \geq 0,$$

this is a contradiction by $|\varepsilon| < \alpha$ and however, we have $\left| \frac{(f*h)(z)}{z} \right| \geq \alpha$. If $g(z) = z - \sum_{n=2}^{\infty} b_n z^n \in \mathcal{N}_\alpha(f)$, then

$$\begin{aligned} \alpha - \left| \frac{(g * h)(z)}{z} \right| &\leq \left| \frac{((f - g) * h)(z)}{z} \right| \leq \sum_{n=2}^{\infty} |a_n - b_n| |c_n| |z^n| \\ &< \sum_{n=2}^{\infty} \frac{\Phi^\eta(\lambda, \mu, n) |n(1 + k) - u_n(k + \gamma)|}{1 - \gamma} |a_n - b_n| \leq \alpha. \end{aligned}$$

□

4. Partial sums of the function class $k - \widetilde{\mathcal{US}}_s^\eta(\lambda, \mu, \gamma, t)$

In this section, applying methods used by Silverman [15] and Silvia [16], we investigate the ratio of a function of the form (1.10) to its sequence of partial sums $f_m(z) = z + \sum_{n=2}^m a_n z^n$.

Theorem 4.1. *If f of the form (1.1) satisfies the condition (2.1), then*

$$(4.1) \quad \operatorname{Re} \left\{ \frac{f(z)}{f_m(z)} \right\} \geq 1 - \frac{1}{\delta_{m+1}}$$

and

$$(4.2) \quad \delta_n \geq \begin{cases} 1, & n = 2, 3, \dots, m \\ \delta_{m+1}, & n = m+1, m+2, \dots \end{cases}$$

where

$$(4.3) \quad \delta_n = \frac{\Phi^\eta(\lambda, \mu, n) |n(k+1) - u_n(k+\gamma)|}{1-\gamma}.$$

The result in (4.1) is sharp for every m , with the extremal function

$$(4.4) \quad f(z) = z + \frac{z^{m+1}}{\delta_{m+1}}.$$

Proof. Define the function $w(z)$, we may write

$$(4.5) \quad \begin{aligned} \frac{1+w(z)}{1-w(z)} &= \delta_{m+1} \left\{ \frac{f(z)}{f_m(z)} - \left(1 - \frac{1}{\delta_{m+1}} \right) \right\} \\ &= \left\{ \frac{1 + \sum_{n=2}^m a_n z^{n-1} + \delta_{m+1} \sum_{n=m+1}^{\infty} a_n z^{n-1}}{1 + \sum_{n=2}^m a_n z^{n-1}} \right\}. \end{aligned}$$

Then, from (4.5) we can obtain

$$w(z) = \frac{\delta_{m+1} \sum_{n=m+1}^{\infty} a_n z^{n-1}}{2 + 2 \sum_{n=2}^m a_n z^{n-1} + \delta_{m+1} \sum_{n=m+1}^{\infty} a_n z^{n-1}}$$

and

$$|w(z)| \leq \frac{\delta_{m+1} \sum_{n=m+1}^{\infty} a_n}{2 - 2 \sum_{n=2}^m a_n - \delta_{m+1} \sum_{n=m+1}^{\infty} a_n}.$$

Now $|w(z)| \leq 1$ if

$$2\delta_{m+1} \sum_{n=m+1}^{\infty} a_n \leq 2 - 2 \sum_{n=2}^m a_n,$$

which is equivalent to

$$(4.6) \quad \sum_{n=2}^m a_n + \delta_{m+1} \sum_{n=m+1}^{\infty} a_n \leq 1.$$

It suffices to show that the left hand side of (4.6) is bounded above by $\sum_{n=2}^{\infty} \delta_n a_n$, which is equivalent to

$$\sum_{n=2}^m (\delta_n - 1) a_n + \sum_{n=m+1}^{\infty} (\delta_n - \delta_{m+1}) a_n \geq 0.$$

To see that the function given by (4.4) gives the sharp result, we observe that for $z = re^{i\pi/n}$,

$$(4.7) \quad \frac{f(z)}{f_m(z)} = 1 + \frac{z^m}{\delta_{m+1}}.$$

Taking $z \rightarrow 1^-$, we have

$$\frac{f(z)}{f_m(z)} = 1 - \frac{1}{\delta_{m+1}}.$$

This completes the proof of Theorem 4.1. □

We next determine bounds for $f_m(z)/f(z)$.

Theorem 4.2. *If f of the form (1.1) satisfies the condition (2.1), then*

$$(4.8) \quad \operatorname{Re} \left\{ \frac{f_m(z)}{f(z)} \right\} \geq \frac{\delta_{m+1}}{1 + \delta_{m+1}}.$$

The result is sharp with the function given by (4.4).

Proof. We may write

$$(4.9) \quad \begin{aligned} \frac{1+w(z)}{1-w(z)} &= (1 + \delta_{m+1}) \left\{ \frac{f_m(z)}{f(z)} - \frac{\delta_{m+1}}{1 + \delta_{m+1}} \right\} \\ &= \left\{ \frac{1 + \sum_{n=2}^m a_n z^{n-1} - \delta_{m+1} \sum_{n=m+1}^{\infty} a_n z^{n-1}}{1 + \sum_{n=2}^{\infty} a_n z^{n-1}} \right\}, \end{aligned}$$

where

$$w(z) = \frac{(1 + \delta_{m+1}) \sum_{n=m+1}^{\infty} a_n z^{n-1}}{- \left(2 + 2 \sum_{n=2}^m a_n z^{n-1} - (1 - \delta_{m+1}) \sum_{n=m+1}^{\infty} a_n z^{n-1} \right)},$$

and

$$(4.10) \quad |w(z)| \leq \frac{(1 + \delta_{m+1}) \sum_{n=m+1}^{\infty} a_n}{2 - 2 \sum_{n=2}^m a_n + (1 - \delta_{m+1}) \sum_{n=m+1}^{\infty} a_n} \leq 1.$$

This last inequality is equivalent to

$$(4.11) \quad \sum_{n=2}^m a_n + \delta_{m+1} \sum_{n=m+1}^{\infty} a_n \leq 1.$$

It suffices to show that the left hand side of (4.11) is bounded above by $\sum_{n=2}^{\infty} \delta_n a_n$, which is equivalent to

$$\sum_{n=2}^m (\delta_n - 1) a_n + \sum_{n=m+1}^{\infty} (\delta_n - \delta_{m+1}) a_n \geq 0.$$

This completes the proof of Theorem 4.2. $\square$

We next turn to ratios involving derivatives.

Theorem 4.3. *If f of the form (1.1) satisfies the condition (2.1), then*

$$(4.12) \quad \operatorname{Re} \left\{ \frac{f'(z)}{f'_m(z)} \right\} \geq 1 - \frac{m+1}{\delta_{m+1}},$$

$$(4.13) \quad \operatorname{Re} \left\{ \frac{f'_m(z)}{f'(z)} \right\} \geq \frac{\delta_{m+1}}{1+m+\delta_{m+1}}$$

where

$$\delta_n \geq \begin{cases} 1, & n = 1, 2, 3, \dots, m \\ n^{\frac{\delta_{m+1}}{m+1}}, & n = m+1, m+2, \dots \end{cases}$$

and δ_n is defined by (4.3). The estimates in (4.12) and (4.13) are sharp with the extremal function given by (4.4).

Proof. Firstly, we will give proof of (4.12). We write

$$\begin{aligned} \frac{1+w(z)}{1-w(z)} &= \delta_{m+1} \left\{ \frac{f'(z)}{f'_m(z)} - \left(1 - \frac{1+m}{\delta_{m+1}} \right) \right\} \\ &= \left\{ \frac{1 + \sum_{n=2}^m n a_n z^{n-1} + \frac{\delta_{m+1}}{m+1} \sum_{n=m+1}^{\infty} n a_n z^{n-1}}{1 + \sum_{n=2}^m a_n z^{n-1}} \right\}, \end{aligned}$$

where

$$w(z) = \frac{\frac{\delta_{m+1}}{m+1} \sum_{n=m+1}^{\infty} n a_n z^{n-1}}{2 + 2 \sum_{n=2}^m n a_n z^{n-1} + \frac{\delta_{m+1}}{m+1} \sum_{n=m+1}^{\infty} n a_n z^{n-1}}$$

and

$$|w(z)| \leq \frac{\frac{\delta_{m+1}}{m+1} \sum_{n=m+1}^{\infty} n a_n}{2 - 2 \sum_{n=2}^m n a_n - \frac{\delta_{m+1}}{m+1} \sum_{n=m+1}^{\infty} n a_n}.$$

Now $|w(z)| \leq 1$ if and only if

$$(4.14) \quad \sum_{n=2}^m n a_n + \frac{\delta_{m+1}}{m+1} \sum_{n=m+1}^{\infty} n a_n \leq 1,$$

since the left hand side of (4.14) is bounded above by $\sum_{n=2}^{\infty} \delta_n a_n$.

The proof of (4.13) follows the pattern of that in Theorem (4.2).

This completes the proof of Theorem 4.3. $\square$

REFERENCES

- [1] Al-Oboudi, F. M. *On univalent functions defined by a generalized Salagean operator*, Int. J. Math. Math. Sci. **27**, 1429-1436, 2004.
- [2] Altıntaş, O. and Owa, S. *Neighborhoods of certain analytic functions with negative coefficients*, Int. J. Math. and Math. Sci. **19**, 797-800, 1996.
- [3] Altıntaş, O., Özkan, E. and Srivastava, H. M. *Neighborhoods of a class of analytic functions with negative coefficients*, Appl. Math. Let. **13**, 63-67, 2000.
- [4] Cataş, A. *Neighborhoods of a certain class of analytic functions with negative coefficients*, Banach J. Math. Anal., **3** (1), No. 1, 111-121, 2009.
- [5] Deniz, E. and Orhan, H. *The Fekete Szegő problem for a generalized subclass of analytic functions*, Kyungpook Math. J. **50**, 37-47, 2010.
- [6] Deniz, E. and Orhan, H. *Some properties of certain subclasses of analytic functions with negative coefficients by using generalized Ruscheweyh derivative operator*, Czechoslovak Math. J., **60** (135), 699-713, 2010.
- [7] Goodman, A. W. *Univalent functions and nonanalytic curves*, Proc. Amer. Math. Soc., **8**, 598-601, 1957.
- [8] Goodman, A. W. *On uniformly starlike functions*, J. Math. Anal. Appl. **155**, 364-370, 1991.
- [9] Orhan, H. *On neighborhoods of analytic functions defined by using hadamard product*, Novi Sad J. Math., **37** (1), 17-25, 2007.
- [10] Owa, S., Sekine, T. and Yamakawa, R. *On Sakaguchi type functions*, Appl. Math. Comput. **187**, 356-361, 2007.
- [11] Răducanu, D. and Orhan, H. *Subclasses of analytic functions defined by a generalized differential operator*, Int. Journal of Math. Analysis **4** (1), 1-15, 2010.
- [12] Ruscheweyh, S. *Neighborhoods of univalent functions*, Proc. Amer. Math. Soc., **81** (4), 521-527, 1981.
- [13] Sakaguchi, K. *On a certain univalent mapping*, J. Math. Soc. Japan **11**, 72-75, 1959.
- [14] Sălăgean, G. S. *Subclasses of univalent functions*, Complex analysis-Proc. 5th Rom.-Finn. Semin., Bucharest 1981, Part 1, Lect. Notes Math. **1013**, 362-372, 1983.
- [15] Silverman H., *Partial sums of starlike and convex functions*, J. Math. Anal. Appl., **209**, 221-227, 1997.
- [16] Silvia E. M., *Partial sums of convex functions of order $\mathbb{R}$* , Houston J. Math., **11** no. 3, 397-404, 1985.
- [17] Srivastava, H. M. and Aouf, M. K. *A certain fractional derivative operator and its applications to a new class of analytic and multivalent functions with negative coefficients. I and II*, J. Math. Anal. Appl., **171**, 1-13, 1992; J. Math. Anal. Appl., **192**, 673-688, 1995.
- [18] Srivastava, H. M. and Owa, S. (Eds.), *Univalent Functions, Fractional Calculus, and their Applications* (Halsted Press (Ellis Horwood Limited, Chichester), John Wiley and Sons, place State New York, 1989).
- [19] Srivastava H.M. and Owa S. (Editors), *Current Topics in Analytic Function Theory*, World Scientific Publishing Company, Singapore, New Jersey, London, Hong Kong, 1992.

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE, ATATURK UNIVERSITY, ERZURUM, 25240, TURKEY.

E-mail address: mcaglar25@gmail.com, horhan607@gmail.com